

Zoochemical Composition of Selected Sea Stars Collected from the Coastal Waters of Carmen, Agusan Del Norte, Philippines

Angelo Mark P. Walag*, Romeo M. Del Rosario, Oliva P. Canencia

Department of Science Education, University of Science and Technology of Southern Philippines, Cagayan De Oro City, PHILIPPINES.

Submission Date: 22-05-2019; Revision Date: 17-07-2019; Accepted Date: 28-08-2019

ABSTRACT

Background: The marine ecosystem is home to 34 of 36 phyla in the animal kingdom. It is for this reason that substantial amount of marine-derived natural products with potential medicinal and therapeutic applications have been described, isolated and characterized. Majority of the invertebrate phyla have been accounted for, but few studies have been made on Philippine invertebrates. **Aim:** The goal of this research is to determine the zoochemical composition selected sea stars collected from the coastal waters of Carmen, Agusan del Norte, Philippines. **Methods:** Standard methods were employed to determine the zoochemical composition of *Linckia laevigata*, *Protoreaster nodosus* and *Acanthaster planci* ground samples. **Results:** All sea stars showed presence of alkaloids, flavonoids, saponins, triterpenoids and cardiac glycosides while no presence of coumarins and phenolics and tannins were detected. The presence of these zoochemicals in sea stars was confirmed by the available literature although differences were noted particularly on the absence of phenolics in this study. Based on this study, the selected stars appeared to be rich in saponins and steroids which are known to possess biological activity that could be used to develop drugs. **Recommendation:** The other metabolites can also be exploited in the developing marine-derived drugs. Further analyses can be done especially in isolating these compounds and structurally elucidating them. Moreover, the isolated metabolites can also be tested individually for its biological activities. It is also suggested that more studies be conducted on these organisms to further understand their biochemistry which can be utilized before they become endangered or more so, extinct.

Key words: Alkaloids, Phenolics, Saponins, Sea stars, Natural Products Chemistry.

Correspondence:

Prof. Angelo Mark P. Walag,

Department of Science Education, University of Science and Technology of Southern Philippines, C.M. Recto Ave, Lapasan, Cagayan De Oro City-9000, PHILIPPINES.

Phone no: +63 9177000485

Email: walag.angelo@gmail.com

INTRODUCTION

The oceans is the most diverse ecosystem with 34 of the 36 phyla represented.^[1] It is with this reason that a substantial number of marine natural products with potential medicinal and therapeutic applications have been described, isolated, and characterized. Of the 34 phyla present in the oceans, marine invertebrates contain metabolites of unprecedented diversity of molecu-

lar structures and activities.^[2] Various laborious efforts have been made to study the different metabolites present in these organisms that might soon lead to further drug development. These marine invertebrates that have been accounted for belong to a wide range phyla cutting across Porifera, Annelids, Coelenterates (Cnidaria and Ctenophora), Mollusks, Echinoderms and etc.^[1,3,4] In the Philippines, efforts have been made to study secondary metabolites and bioactivities of these invertebrates like ascidians,^[5] corals,^[6] sponges,^[7-11] sea stars, sea cucumbers, and brittle stars,^[12] mollusks.^[13] Although, several studies have been made on Philippine invertebrates, much of the efforts have been focused on sponges and other sessile organisms which are believed to possess diverse chemical defenses. Several phyla are yet to be studied that may possess lead compounds for future

SCAN QR CODE TO VIEW ONLINE



www.ajbls.com

DOI :
10.5530/ajbls.2019.8.10

drug development studies. Moreover, various issues on water quality of both marine and freshwater have been noted that may affect abundance and diversity of the aquatic life.^[14] Several studies were also conducted to evaluate the abundance and diversity of echinoderm species in Mindanao and have noted a decline in their population.^[15-21]

Of the different phyla mentioned, echinoderms are spread worldwide and known to have more than 7000 species divided into five classes.^[22] These echinoderms are also known as producers of bioactive metabolites.^[23] They are also reported to have ability to produce chemical defenses against possible predators although recent studies are rare to find.^[23] Although, this phylum has been found to possess various bioactive secondary metabolites, much of the studies conducted on echinoderms in the Philippines come from and are limited on traditional fields like ecology, biogeography, effects of pollution, and conservation of species. This gap in the literature, particularly in the study of secondary metabolites from echinoderms in the Philippines, further emphasizes the need to exert effort in unravelling the diverse chemical world of secondary metabolites before these animals become endangered due to the compounding effects of climate change, pollution, exploitation, and etc. The identification, structure elucidation, and characterization of secondary metabolites and their ecological role have led to the discovery of lead compounds with relevant pharmacological properties.^[24]

Sea stars are invertebrates which belongs to class Asterozoa, phylum Echinodermata which are often found in marine environments although some are found in brackish waters.^[25] These organisms produce unique natural products which has been found to be in demand in the fields of medicine.^[26] In fact, under the phylum Echinodermata, three classes (Holothuria, Asterozoa, and Crinozoa) have been noted to produce elaborate chemical defenses against predation.^[27] Furthermore, the secondary metabolites they produce are an integral part of their survival strategy to protect them against predators and parasites. These compounds sea stars produce like glycosides, sterols, cerobrosides, peptides, and gangliosides possess a wide range of bioactivities like being cytotoxic, antimicrobial, expectorant, analgesic, hemolytic, antioxidant and antitussive.^[28-31] Sea stars in the Philippines has only been studied in terms of bioactivity against brine shrimp and certain strains of bacteria^[12,32] and no literature yet available on the secondary metabolite screening. Thus, the main aim of this study is to determine the zoochemicals present in selected sea stars from Goso-on and Vinapor, Carmen, Agusan del Norte, Philippines.

MATERIALS AND METHODS

Study Area

The animal materials required for this study was collected from the intertidal zone up to shallow waters (0-5m) of Barangay Goso-on, Carmen, Agusan del Norte, Philippines. The map of the collection site is shown in Figure 1. The Municipality of Carmen is located in the province of Agusan del Norte of CARAGA Region or Region XIII. It is strategically located along the Western Agusan Corregidor, surrounded by the Butuan Bay in the north, Buenavista in the south, Nasipit in the east, and Misamis Oriental in the west.

Collection and identification of sea star specimens

A field reconnaissance was conducted to evaluate presence and abundance of marine sea stars selected for this study. Preliminary identification of specimens was conducted based on the field guide developed by Schoppe (2000) and the World Registry of Marine Species (WoRMS) using morphological characteristics. Specimens that were identified were photographed and then returned back to its natural habitat. One representative per specimen was collected and stored in polypropylene bags for further identification. Final verification and confirmation of species identification was through the use of collected and preserved specimens that were brought to the laboratory. Photographs of specimens were also sent to the Institute of Environmental and



Figure 1: Collection site in Barangay Goso-on, Carmen, Agusan del Norte.

Marine Science, Silliman University for the confirmation of the identification. The sea stars collected were *L. laevigata*, *P. nodosus*, and *A. planici*. A collection of aquatic samples using non-destructive method. Gratuitous Permit (GP-2019-01) was also acquired from the Bureau of Fisheries and Aquatic Resources – Caraga Region prior sampling and reconnaissance.

Fresh samples were collected during the lowest low tide of the month of April 2019 from the intertidal zone up to shallow (3-5 meters deep) parts of the marine environment in Barangays Goso-on and Vinapor, Carmen, Agusan del Norte. Mature (>5 inches across) sea stars that were collected were washed with marine water to remove dirt and sand and were then placed in styrobox with ice and marine water for preservation. All collected samples were brought to Cagayan de Oro City for storage and further sample processing.

Sample Preparation

The specimens were sorted in polypropylene bags filled with marine water. One bag was used for each species to avoid contamination. Specimens were brought to the chemistry laboratory and reduced to smaller size using a clean pruning shear. Samples were subjected to drying using oven method blanketed with nitrogen gas to minimize oxidation. Samples were homogenized separately using a mechanized grinder available at the Northern Mindanao Food Innovation Center. The sample material were weighed and transferred to pre-weighed Ziploc bags. Each sample bag was labelled with its scientific name and a specific sample number.

Zoochemical Analyses

Test for Alkaloids. Presence of alkaloids was determined qualitatively using the Wagner's Test.

1. Wagner's Test. 50 mg (0.05 g) of the sample was stirred with 2 mL dilute HCl (2N) and filtered. The filtrate (2 mL) was added with few drops of Wagner's reagent (1.27 g iodine and 2 g potassium iodide dissolved in 5 mL of water and made up to 100 mL with distilled water). The samples were scored positive on the basis of turbidity or precipitation.

Tests for Coumarin. Presence of coumarins was determined using NaOH test and Ammonium hydroxide-UV light test.

1. NaOH Test. 0.05g of moistened dry powdered sample was placed in a test tube. The test tube was covered with filter paper soaked in dilute NaOH (10%). The mixture was kept in water bath and after some time, the filter paper was exposed to

UV light. A yellowish-green fluorescence indicates presence of coumarin.^[34]

2. Ammonium hydroxide-UV light test. A moistened dry powdered sample (0.05 g) was placed in a test tube and was covered with filter paper soaked in dilute ammonium hydroxide (10%) and was kept in water bath. After some time, the filter paper was exposed to UV light. An intense fluorescence indicates presence of coumarin.

Tests for Flavonoids. The presence of flavonoids was determined using Shinoda's test and ammonia test.

1. Shinoda's Test. 400 mg (0.4 g) sample was extracted with 10 mL ethanol was then filtered. 1 mL filtrate was added with magnesium ribbon and conc. HCl (5 drops) and allowed to react. A pink or red color indicated the presence of flavonoids.
2. Ammonia test. 30 ml of ethyl acetate was added to about 0.6 g of the powdered sample and was heated on a water bath for 3 minutes. The mixture was cooled, filtered and 4 mL of filtrate was shaken with 1 mL of dilute ammonia solution (10%). The layers are allowed to separate and the yellow color in the ammonia layer indicates the presence of flavonoids

Test for Phenolics and Tannins. The presence of phenolics and tannins was determined using ferric chloride test.

1. Ferric chloride test. 500 mg (0.5 g) sample was boiled in 30 mL distilled water and filtered. 2 mL filtrate was mixed with 1 mL 5% FeCl₃. Formation of blue, green, or violet color indicated the presence of phenolic and tannin compounds.

Tests for Saponins. The presence of saponins was indicated using foam test and olive oil test.

1. Foam test. 1 g powdered sample was boiled in 10 mL distilled water for 15 min. After cooling, the extract was shaken vigorously to record froth formation. Presence of honeycomb froth above the surface after 30 mins indicates the presence of saponins.
2. Olive oil test. About 2 g of the powdered sample will be boiled in 20 mL of water in a water bath and will then be filtered. 10 mL of the filtrate will be mixed with 5 mL distilled water and shaken vigorously to form a stable persistent froth. The froth will then be mixed with 3 drops of olive oil, shaken vigorously, and then observed for the formation of emulsions.

Test for Triterpenoids. The presence of triterpenoids was determined using Salkowski test.

1. Salkowski test. 0.6 g of the sample was extracted in 30 mL chloroform, and filtered. 2 mL of the

filtrate was treated with few (3-4) drops of concentrated sulfuric acid, shaken and allowed to stand. Presence of triterpenoids was indicated by the appearance of yellow color in the lower layer.

Tests for Steroids. The presence of steroids was indicated using the Liebermann-Burchard test and Salkowski test.

1. Liebermann-Burchard test. 0.6 g of the sample was extracted in 30 mL chloroform, and filtered. 2 mL of the filtrate was added with 2 mL acetic anhydride and 1 mL conc. sulfuric acid and allowed to react. A blue-green ring indicated the presence of steroids
2. Salkowski test. 0.6 g of the sample was extracted in 30 mL chloroform, and filtered. 2 mL of the filtrate was treated with few (3-4) drops of concentrated sulfuric acid, shaken and allowed to stand. Presence of steroids was indicated by the appearance of red color in the lower layer.

Test for Cardiac glycoside. The presence of cardiac glycosides was determined using the test for Keller-Kiliani Test.

1. Keller-Kiliani Test. 500 mg sample was boiled in 30 mL distilled water and filtered. 2 mL of the filtrate was added with 1 mL glacial acetic acid and 1 mL 5% FeCl₃ + 1 mL conc. sulfuric acid. A green-blue color indicated the presence of glycosides.

RESULTS

Zoochemical Composition

Alkaloids are commonly defined as any substance which contain one or more nitrogen atoms usually as part of a cyclic system.^[35] In plants, alkaloids are known to function as protective agents against predators,

growth regulators, and as waste products in metabolisms.^[36] Alkaloids were found to be present in all three sea stars as indicated in the Wagner's Test result which was scored based on the presence of turbidity or precipitation as summarized in Table 1. Coumarin is the parent compound of a large class of naturally occurring phenolic compounds and have been found to be widely distributed to various plant families while only few were found in animals.^[37,38] Coumarins were found to be absent in all sea stars investigated in this research. Flavonoids are known in plants as pigments, synthesized from phenylalanine, which exudes brilliant fluorescence when excited with UV light.^[39] The production of flavonoids in living organisms has been believed to be a response against fungal parasites, predation, pathogens, and oxidative cell injury.^[40] The presence of flavonoids in the selected sea star samples are presented in Table 1. All three sea stars possess flavonoids as indicated by Shinoda's Test and Ammonia Test although differing in terms of their intensity. All sea stars investigated were found to possess flavonoids in both tests conducted. Phenolics and tannins, on the other hand, were found to be absent in all sea stars. Saponins, triterpenoids, steroids, and cardiac glycosides were also found to be present in all sea stars ranging from trace presence to strong presence as summarized in Table 1.

DISCUSSION

Trace to moderate presence was observed in the three samples which coincide with the finding that echinoderms were the second least nitrogen-bearer compared to different marine invertebrates.^[41] Furthermore, Baker reported that the three major producers of alkaloids, in proportion, in the marine ecosystem are organisms

Table 1: Zoochemicals present in selected sea stars using varied tests.

Zoochemical	Test	<i>L. laevigata</i>	<i>P. nodosus</i>	<i>A. planci</i>
Alkaloids	Wagner's Test	++	+	+
	NaOH Test	-	-	-
Coumarin	NH ₄ OH-UV Light Test	-	-	-
	Shinoda's Test	++	+	+
Flavonoids	Ammonia Test	+	++	++
	Ferric chloride Test	-	-	-
Phenolics and Tannins	Foam Test	++	++	++
	Olive Oil Test	++	++	++
Triterpenoids	Salkowski's Test	+	++	+
	Liebermann-Burchard Test	++	+++	++
Steroids	Salkowski's Test	+	++	++
	Keller-Kiliani Test	++	+	+

Legend: +++ strong presence, ++ moderate presence, + trace presence and - absence.

under taxa of tunicates, worms, and bryozoan. Sponges were also reported to possess the most rich source of alkaloids in the marine ecosystem.^[42] It was also noted that majority of the nitrogenous substances found in marine organisms are halogenated (25%), sulfur-bearing in one of several oxidation states (13%), polypeptides (8%), and possess N-O bond (7%).

Alkaloids were also isolated from sea stars *Lethasterias nanimensis chelifera*,^[30] *Perknaster fuscus antarcticus* (Antarctic sea star) which is considered to be the most ichthyotoxic echinoderm in the Antarctic,^[43] and *Dermasterias imbricata*^[44] Several alkaloidal substances were also isolated from *Pentaceraster chinensis*,^[45] which possesses antimicrobial activity against Gram-positive and -negative bacteria, *Fromia monilis* and *Celerini heffermani*, which also possess cytotoxic activities.^[46] These alkaloids, together with many other alkaloids, possess toxic effects by mimicking structurally and biosynthetically related neurotransmitters.^[47] This further highlights that the production of alkaloids in sea stars may be attributed to its chemical defense nature as suggested by Waller and Nowacki (1978).

The moderate presence of alkaloids in *L. laevigata* suggests the need for further protection against possible predators due to the absence of thorns in its body walls. *P. nodosus* and *A. planci* possess a thick body wall and an elaborate system of thorns, respectively, which may be the reason of low alkaloid presence. Alkaloids are known to taste bitter in plants, which may also deter predators in marine ecosystems. Furthermore, it was also observed that both *P. nodosus* and *A. planci* specimens collected were relatively larger than *L. laevigata*. In marine organisms, tradeoff between growth and development and variation in defenses has been well observed in the Optimal Defense Theory first proposed by Rhoades (1979).

In marine ecosystems, two isocoumarins were found present in sponge-associated fungus *Aspergillus similanensis*^[48] while one isocoumarin was isolated from the marine-derived fungus *Aspergillus flavus* (from a prawn species *Penaeus vannamei*).^[49] Another species of *Aspergillus*, a marine-derived fungus, yielded a new dihydroxyisocoumarin from the ethyl acetate extract which showed inhibition of LPS-induced nitric oxide and prostaglandin E₂.^[50] Another isocoumarin compound was isolated from marine-derived fungus associated with seagrass.^[51,52] This shows that most coumarins in marine ecosystems are derived from associated fungus thus supports results of this study particularly in the absence of coumarins in the three sea star species.

The presence of flavonoids as coloring pigments in sea stars may be an elaborate defense mechanism to serve

as warning against potential predators. This mechanism of possessing aposematic coloration has been well observed in small and obscure organisms.^[53] Furthermore, organisms under the phylum Echinodermata, includes 6000 species possessing conspicuously colored parts like sea cucumbers, brittle stars, and sea stars. Moreover, asteroids (sea stars) are believed to have evolved with chemical defenses as an in part of their survival strategies.^[26]

L. laevigata are sea stars characterized by uniformly blue color due to a blue, grey, pink, or purple pigment.^[33] In the Philippines, *L. laevigata* species have been found to be in shades of blue^[15,16] which is due to a pigment called linkiacyanin.^[54] This linkicyanin is a compound with varied pigments that reflect their availability in terms of the animal's diet or metabolic activities. The role of this coloration in the skin of sea stars has been believed to be an adaptive camouflage rather than as light receptors.^[54] This camouflage coloration in sea stars allows them to hide against possible predators. Similarly, *A. planci*'s coloration varies from red to orange which also reflects its variability as an attribute to the animal's diet.^[33] Red and orange are the most frequent colors among phylum Echinodermata, as exhibited by both *A. planci* and *P. nodosus*, which are due to varied combinations of carotenoids, carotenoproteins, flavins, melanins, quinones, and porphyrins.^[55]

Phenolic compounds are naturally occurring antioxidants found in plants and other marine invertebrates particularly sea cucumbers and sea urchins.^[31] Furthermore, sea cucumbers were found to possess high phenolic content in their digestive tract while low phenolic content in their gonads^[56] *Holothuria scabra*, *Holothuria leucospilota*, and *Stichopus chloronatus* from Malaysia^[57] *Strongylocentrotus droebachiensis*,^[58] and *Curcumaria frondosa*.^[56] These phenolic compounds collected from these sea stars also showed antioxidant activity.

Sea stars were also found to possess phenolic compounds. Reasonable amounts of phenolics were determined in three different extracts from *L. maculata* which can be a good source of antioxidants.^[31] Pharmacological and chemical properties of phenolic compounds were also evaluated in two sea star species (*Astropecten irregularis* and *Luidia sarsii*) and one brittle star (*Ophiura albida*).^[59] Phenolic compounds were also present in organic extract of brittle star *Ophiocoma erinaceus* which showed antioxidant anti-inflammatory properties.^[60] Phenolics and tannins showed negative in all three star species although present in many studies on echinoderms, even in *A. planci* where phenolic compounds were identified.^[61] and showed antioxidant and anticancer activities. The absence may be attributed to the fact

that majority of the phenolics identified in the literature came from the pyloric caeca of these organisms and the material analyzed for this study is only the body wall.

Saponins are naturally occurring non-volatile surface-active glycosides mainly produced by plants but also lower marine animals and bacteria.^[62,63] Saponins in sea stars generally play important roles in digestion, reproduction, and chemical signaling.^[64] Saponins were moderately present in all sea star samples which is not surprising since a good amount of literature on saponin isolation in sea stars and other echinoderm species are available. To note, saponins were isolated in two different sea cucumbers *H. lessona*^[65] and *H. moebii*^[66] where the latter showed *in vitro* cytotoxicity against tumor growth in mice. Not only sea cucumber possess saponins but also sea stars, in fact, among echinoderms, sea stars and sea cucumbers possess saponins that is generally responsible for their toxicity.^[67]

Glycosylated steroids produced by sea stars bear one to six sugar groups and are collectively called saponins because of their soap-like tendencies to form emulsions with water. Saponin groups commonly found in sea stars are often sulfonylated^[25] which are referred to as asterosaponins. Several sea stars around the world have been a source of a handful of asterosaponins. Two new asterosaponins (luidiaquinoside and psilasteroside) were isolated from *Luidia quinaria* and *Psilaster cassiope* which showed marginal *in vitro* cytotoxicity against rat basophilic leukemia cells.^[68] Sulfated steroidal glycosides together with three known asterosaponins were also isolated from *Culcita novaeguineae* which all showed moderate cytotoxicity against cancer cell lines.^[69] Six new asterosaponins were also isolated from *Leptasterias ochotensis* which showed slight to moderate cytotoxicity against cancer cell lines.^[70] Anti-inflammatory asterosaponins were also isolated from the ethanolic extract of an edible Vietnamese sea star *Astropecten monacanthus*.^[71]

The saponins detected in this study can be confirmed by the different saponin isolation and structural elucidation available in the literature. Five saponin compounds were also isolated from *L. laevigata*, where four of which were known (thornasteroside, marthasteroside, ophidianside, and maculatoside) and the other one was named as laevigatoside, where all of which were found to be steroidal glycoside sulfates.^[72] A novel asterosaponin nodoside were also isolated from the sea stars *A. planici* and *L. laevigata*.^[73] Another steroidal glycoside sulfate compound was isolated from the sea star *P. nodosus* and was further structurally elucidated.^[74] Moreover, in a more recent study, polyhydroxylated sterols isolated from *P. nodosus* showed anti-inflammatory activities.^[75] This

abundant literature on saponin presence in sea stars goes to show that asteroids are good source of novel saponin compounds. The toxic properties of sea stars are believed to be associated with compounds similar to plant saponins.^[61,67] Furthermore, it is because of the toxic nature of saponins that it is probable that saponins act as chemical defense against infectious fungi, protists, parasites, and predators.^[76] This could be the reason for toxic nature of the spines of *A. planici* which showed positive presence of saponins. In addition, the butanolic fraction extract from *A. planici* contained cytotoxic asterosaponins.^[61]

Triterpenoids are considered to be a ubiquitous non-steroidal secondary metabolite in both marine and terrestrial fauna.^[77] Triterpene glycosides have been initially believed to be present and abundant only in sea cucumbers^[78] in the marine fauna but later were found to be present in marine sponges.^[79] In a more recent review, it was found out that triterpenoids are the most abundant secondary metabolite present in marine organisms (sponges, sea cucumbers, marine algae, marine-derived fungi).^[80] In addition, four triterpenoids were also described and isolated from a marine sponge *Jaspis stellifera* in a more recent study.^[81] In sea cucumbers, the triterpene glycosides are synthesized in the skin and are ejected when the animal is disturbed. This behavior is believed to be a defense mechanism holothuroid species possess.^[82]

The abundance of triterpenoids in the marine life can be confirmed by the presence of this compound in all three sea star samples as indicated in the Salkowski's Test. Although present in the specimens tested, it was suggested that the saponins present in sea cucumbers are triterpenoid glycosides while in sea stars are steroidal glycosides.^[67] Furthermore, it was also pointed out agree that sea cucumbers are producers of triterpenoid glycosides while sea stars are steroidal glycosides.^[76] This could have been the reason for the non-availability of literature on the sea star triterpenoid compounds. Moreover, the presence of oligoglycosides in both asteroids and holothuroids suggests the phylogenetic closeness of both taxa.

Steroids are compounds which possess a fundamental structure of four fused carbon rings commonly known as steroidal nucleus^[83] Sea stars are known to be excellent producers of secondary metabolites particularly steroids.^[25] Moreover, the vast majority (>80%) of the secondary metabolites discovered in sea stars are steroids.^[31] Unlike other echinoderms, sea stars are known to produce a wide variety of oxygenated steroids compounds like polyhydroxysteroids, polyhydroxysteroid mono- and biosides, and toxic steroid oligoglycosides.

^[84] Several sea stars have been found to possess steroidal compounds where majority of which exhibits anti-neoplastic, antiviral, anti-inflammatory, and ichthyotoxic activities.^[85] Three sulfated steroidal pentaglycosides have been isolated from the sea star *C. novaeguineae* which all possessed marginal cytotoxic activity.^[86]

As shown in Table 1, all three sea stars samples showed moderate to strong presence of steroids. This confirms the idea that sea stars are considered to be richest source of polyhydroxy steroids in the marine ecosystem.^[87] Furthermore, steroids were identified in *P. nodosus* and were found to possess the most common steroids in sea stars.^[88] Three more steroids were further isolated from the same sea star which was also found to be common in other sea stars.^[73] Cytotoxic steroids were also isolated first from the same sea star which was later called cytotoxic nodoside.^[87] Four steroids were also isolated from *A. planici* which also confirms the presence of steroids of this same species in this study.^[89] Several studies also reported presence of steroidal glycosides present in *A. planici*.^[90,91]

Steroid glycosides were also isolated from *L. laevigata* which confirms the results of this study. Two new steroids were isolated from this sea star and were found to possess two monosaccharide units.^[92] Three more novel steroids were also isolated from the same sea star were found to possess two monosaccharide units.^[93] These steroids isolated were described to be members of neurogenic compounds.

Cardiac glycosides or cardiotoxic glycosides possess a steroidal structure with an unsaturated lactone ring.^[94] This metabolite is present in a limited number of plant families and is clinically used to treat congestive heart failure.^[95,96] In marine ecosystems, cardiac glycosides together with steroid saponins consists the toxin called holothurin. This toxin produced by sea cucumbers have been found to cause skin papular eruptions and eye irritation.^[97] Ingestion of this holothurin containing potent cardiac glycoside has also been noted to cause death.^[98] This toxin produced by sea cucumbers might be related to cardiac glycosides detected in this study. These cardiac glycosides detected might serve as defense mechanism of sea stars against predators and parasites.

CONCLUSION

The presence of alkaloids in the three sea stars species coincides with the finding that echinoderms were the second least nitrogen-bearer compared to different marine invertebrates. The results also suggest that the presence of alkaloids in sea stars serves the role as protective agent against predators, as growth regulators,

and as waste products of metabolism. This also suggests that differences in alkaloid presence in sea stars can be attributed to presence of potential predators. The absence of coumarins in sea stars was not surprising since majority of the coumarins isolated from marine ecosystems come from fungi associated with sponges, prawns, and sea grasses. This further suggests that sea stars do not produce coumarins as part of their defense mechanism. Flavonoids in sea stars serve as their coloring pigments which is part of their elaborate defense mechanism, serving as an advanced warning against predators. Phylum Echinodermata is known to possess conspicuously colored parts to deter potential predators. The differences of flavonoid presence can be attributed to the color of this marine organism. The more colorful, the more flavonoid compounds can be expected. The absence of phenolics in this study may be attributed to the fact that majority of the phenolics identified in the literature came from the pyloric caeca of these organisms and the material analyzed for this study is only the body wall. Saponins are generally present in sea stars which serve as chemical defense against infectious fungi, protists, and parasites, and predators. The presence of saponins in sea stars also supports the idea this metabolite possess toxic properties similar to saponin counterparts in plants. Triterpenoids are generally absent in sea stars since they are known to be producers of steroidal glycosides while the major producers of echinoderm triterpenoids are sea cucumbers. The triterpene glycosides present in sea stars in this study is believed to be part a defense mechanism of these organisms similar to those of holothuroid species. Sea stars are considered to be richest source of polyhydroxy steroids in the marine ecosystem. These saponins collected from these organisms are believe to be part of their defense mechanism as saponins from other sea star showed potent cytotoxic activity against broad range of test subjects. Cardiac glycosides are known to be produced by sea cucumbers as part of the toxin they produce. The cardiac glycosides detected in this study are believed to be part of the elaborate defense mechanism of these organisms against predators and parasites especially in the toxic spines of *A. planici*.

ACKNOWLEDGEMENT

The authors would like to express their immense gratitude to the Bureau of Fisheries and Aquatic Resources – Caraga Region for the issuance of the necessary Gratuitous Permit and all assistance provided for this research. Same gratitude is also expressed to the Agriculture Office of the Municipality of Carmen for the

issuance of a Municipal Clearance in consultation with the Protected Areas Management Board and other relevant offices. The authors would also like to thank Mr. Archie Layaog, Mr. Efren G. Tangon, Ms. Genevieve Garcia, and Ms. Mary Rose Sullano for the assistance provided in the sampling and conduct of the experiments of this research.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

SUMMARY

All sea stars showed presence of alkaloids, flavonoids, saponins, triterpenoids and cardiac glycosides while no presence of coumarins and phenolics and tannins were detected. Moreover, the selected stars appeared to be rich in saponins and steroids which are known to possess biological activity that could be used to develop drugs.

REFERENCES

- Sima P, Vetvicka V. Bioactive substances with anti-neoplastic efficacy from marine invertebrates: Porifera and Coelenterata. *World J Clin Oncol* [Internet]. 2011;2(11):355. Available from: <http://www.wjgnet.com/2218-4333/full/v2/i11/355.htm>
- Thao NP, No JH, Luyen BTT, Yang G, Byun SY, Goo J, et al. Secondary metabolites from vietnamese marine invertebrates with activity against *trypanosoma brucei* and *T. cruzi*. *Molecules*. 2014;19(6):7869–80.
- Blunt JW, Copp BR, Keyzers RA, Munro MHG, Prinsep MR. Marine natural products. *Nat Prod Rep* [Internet]. 2016;33(3):382–431. Available from: <http://xlink.rsc.org/?DOI=C5NP00156K>
- Leal MC, Puga J, Seródio J, Gomes NCM, Calado R. Trends in the discovery of new marine natural products from invertebrates over the last two decades - where and what are we bioprospecting? *PLoS One*. 2012;7(1).
- Perez LJ, Faulkner DJ, Fenical W. Bistratamides E-J, modified cyclic hexapeptides from the Philippines ascidian *Lissoclinum bistratum*. *J Nat Prod*. 2003;66(2):247–50.
- Xio YJ, Su JH, Tseng YJ, Chen BW, Liu W, Sheu JH. Oxygenated eremophilane- and neolemnane-derived sesquiterpenoids from the soft coral *Lemnalia philippinensis*. *Mar Drugs*. 2014;12(8):4495–503.
- Rivera AP, Uy MM. In vitro antioxidant and cytotoxic activities of some marine sponges collected off misamis oriental coast, Philippines. *E-Journal Chem*. 2012;9(1):354–8.
- Tatsukawa M, Punzalan LLC, Magpantay HDS, Villaseñor IM, Concepcion GP, Suwanborirux K, et al. Chemistry of renieramycins. Part 13: Isolation and structure of stabilized renieramycin type derivatives, renieramycins W-Y, from Philippine blue sponge *Xestospongia* sp., pretreated with potassium cyanide. *Tetrahedron*. 2012;68(36):7422–8.
- Lerch ML, Harper MK, Faulkner DJ. Brominated polyacetylenes from the Philippines sponge *Diplastrella* sp. *J Nat Prod*. 2003;66(5):667–70.
- Yamakuma M, Kato H, Matsuo K, El-Desoky AH, Kawabata T, Losung F, et al. 1-Hydroxyethylhalenaquinone: A new proteasome inhibitor from the marine sponge *Xestospongia* SP. *Heterocycles*. 2014;89(11):2605–10.
- Quiao MAD, Uy MM. Pyrimidines from the Philippine Marine Sponge *Aaptos suberitoides*. *Int J Sci Eng Res*. 2013;4(1):1–4.
- Layson RJ, Rodil MCA, Mojica EE, Deocaris CC. Potential Anti-cancer and Anti-bacterial Activities of Philippine Echinoderm Extracts. *J Trop Life Sci*. 2014;4(3):175–81.
- Jimenez EC, Shetty RP, Lirazan M, Rivier J, Walker C, Abogadie FC, et al. Novel excitatory *Conus* peptides define a new conotoxin superfamily. *J Neurochem*. 2003;85(3):610–21.
- Walag AMP, Canencia OP, Fiedler BA. Water Quality: Mindanao Island of the Philippines. In: Fiedler BA, editor. *Translating National Policy to Improve Environmental Conditions Impacting Public Health Through Community Planning* [Internet]. Cham: Springer International Publishing; 2018. p. 219–53. Available from: https://link.springer.com/chapter/10.1007/978-3-319-75361-4_12
- Llacuna MEJ, Walag AMP, Villaluz EA. Diversity and dispersion patterns of echinoderms in Babanlagan, Talisayan, Misamis Oriental, Philippines. *Environ Exp Biol*. 2016;14(4):213–7.
- Walag AMP, Layaog AG, Garcia GU. Survey of echinoderms in the intertidal zone of Goso-on and Vinapor, Carmen, Agusan del Norte, Philippines. *Environ Exp Biol*. 2018;16(1):31–8.
- Walag AMP, Canencia MOP. Physico-chemical parameters and macrobenthic invertebrates of the intertidal zone of Gusa, Cagayan de Oro City, Philippines. *AES Bioflux* [Internet]. 2016 Apr;8(1):71–82. Available from: <http://www.aes.bioflux.com.ro/docs/2016.71-82.pdf>
- Medrano MGT. Diversity of macrobenthic invertebrates in the intertidal zone of Brgy. Tagpangahoy, Tubay, Agusan del Norte, Philippines. *Int J Tech Res Appl*. 2015;19(Special Issue):5–9.
- Pitogo KME, Sumin JP, Ortiz AT. Shallow-water Sea Cucumbers (Echinodermata: Holothuroidea) in Sarangani Bay, Mindanao, Philippines with Notes on Their Relative Abundance. *Philipp J Sci*. 2018;147(3):453–61.
- Lacuna MLDG, Merlas GP. Macrobenthic fauna on the rocky intertidal shore of Sultan Naga Dimaporo, Lanao del Norte, Mindanao, Philippines. *Int J Biol Pharmacy, Appl Sci*. 2018;7(9):1675–85.
- Pan M, Bailly N, Conejar J, Dar C, Froese R, Garilao C, et al. Philippine marine biodiversity as assessed by SeaLifeBase: current status. *UPV J Nat Sci*. in press:1–83.
- Dupont S, Ortega-Martínez O, Thorndyke M. Impact of near-future ocean acidification on echinoderms. Vol. 19, *Ecotoxicology*. 2010. p. 449–62.
- Skropeta D, Wei L. Recent advances in deep-sea natural products. *Nat Prod Rep* [Internet]. 2014;31(8):999–1025. Available from: <http://xlink.rsc.org/?DOI=C3NP70118B>
- Mayer AMS, Rodríguez AD, Berlink RGS, Hamann MT. Marine pharmacology in 2005-6: Marine compounds with anthelmintic, antibacterial, anticoagulant, antifungal, anti-inflammatory, antimalarial, antiprotozoal, antituberculosis, and antiviral activities; affecting the cardiovascular, immune and nervous system. *Biochim Biophys Acta - Gen Subj* [Internet]. 2009;1790(5):283–308. Available from: <http://dx.doi.org/10.1016/j.bbagen.2009.03.011>
- Dong G, Xu T, Yang B, Lin X, Zhou X, Yang X, et al. Chemical constituents and bioactivities of starfish. *Chem Biodivers*. 2011;8(5):740–91.
- Sumithra R, Banu N, Deepa Parvathi V. Novel Natural Products from Marine Sea Stars. *Curr Trends Miomedical Eng Biosci*. 2017;2(4):1–5.
- Slattery M. Bioactive compounds from echinoderms: Ecological and evolutionary perspectives. In: *Echinoderms: Durham - Proceedings of the 12th International Echinoderm Conference*. 2010. p. 591–600.
- Bi L, Tian X, Dou F, Hong L, Tang H, Wang S. New antioxidant and antilycogen active triterpenoid saponins from the root bark of *Aralia taibaiensis*. *Fitoterapia*. 2012;83(1):234–40.
- Cheng G, Zhang X, Tang HF, Zhang Y, Zhang XH, Cao WD, et al. Asterosaponin 1, a cytostatic compound from the starfish *Culcita novaeguineae*, functions by inducing apoptosis in human glioblastoma U87MG cells. *J Neurooncol*. 2006;79(3):235–41.
- Kicha AA, Ivanchina N V., Kalinovskiy AI, Dmitrenok PS, Stonik VA. Alkaloidosteroids from the starfish *Lethasterias nanimensis* chelifera. *Tetrahedron Lett*. 2003;44(9):1935–7.
- Suguna A, Bragadeeswaran S, Natarajan E, Mohanraj M. Studies on antioxidant properties of starfish *Luidia maculata*. *J Coast Life Med* [Internet]. 2014;2(9):694–8. Available from: <http://www.jclmm.com/qk/20149/5.pdf>
- Walag AMP, Del Rosario RM. Proximate biochemical composition and brine shrimp lethality assay of selected sea stars from Goso-on and Vinapor, Carmen, Agusan del Norte, Philippines. *Malaysian J Biochem Mol Biol*. 2018;21(3):11–8.
- Schoppe S. Echinoderms of the Philippines: A guide to common shallow water sea stars, brittle stars, sea urchins, sea cucumbers and feather stars.

- Philippines: VISCA-GTZ Program on Applied Tropical Ecology, Visayas State College of Agriculture; 2000. 144 p.
34. Walag AMP, Cepeda ABM, Galenzoga AS, Sambaan SMJ. Initial phytochemical screening of the different parts of *Mansoa alliacea* L. (Garlic Vine). *Int J Biosci* [Internet]. 2017;11(3):227–31. Available from: <http://www.innspub.net/ijb/initial-phytochemical-screening-different-parts-mansoa-alliacea-l-garlic-vine/>
 35. Harborne JB. *Phytochemical Methods: A Guide To Modern Techniques Of Plant Analysis*. Third. Vol. 3, Chapman and Hall. London: Chapman and Hall; 1998. 317 p.
 36. Waller GR, Nowacki EK. The Role of Alkaloids in Plants BT - *Alkaloid Biology and Metabolism in Plants*. In: Waller GR, Nowacki EK, editors. Boston, MA: Springer US; 1978. p. 143–81. Available from: https://doi.org/10.1007/978-1-4684-0772-3_5
 37. Riveiro ME, De Kimpe N, Moglioni a, Vázquez R, Monczor F, Shayo C, et al. Coumarins: old compounds with novel promising therapeutic perspectives. *Curr Med Chem*. 2010;17(13):1325–38.
 38. Soine TO. Naturally occurring coumarins and related physiological activities. *J Pharm Sci* [Internet]. 1964;53(3):231–64. Available from: http://www.journals.cambridge.org/abstract_S0952675714000244
 39. Havsteen BH. The biochemistry and medical significance of the flavonoids. *Pharmacol Ther*. 2002;96(2–3):67–202.
 40. Cook NC, Samman S. Flavonoids—Chemistry, metabolism, cardioprotective effects, and dietary sources. *J Nutr Biochem*. 1996;7(2):66–76.
 41. Iken KB, Baker BJ. Ainigmaptilonenes, sesquiterpenes from the Antarctic gorgonian coral *Ainigmaptilon antarcticus*. *J Nat Prod* [Internet]. 2003;66(6):888–90. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/12828486>
 42. Baker BJ. Chapter Four β -Carboline and isoquinoline alkaloids from marine organisms. *Alkaloids Chem Biol Perspect*. 1996;10(C):357–407.
 43. Kong F, Harper MK, Faulkner DJ. Fuscusine, A Tetrahydroisoquinoline Alkaloid from the Seastar *Perknaster fuscus* Antarcticus. *Nat Prod Lett*. 1992;1(1):71–4.
 44. Ohba M, Nishimura Y, Kato M, Fujii T. Synthetic studies on the starfish alkaloid imbricatine. A chiral synthesis of tri-O-methylimbricatine. *Tetrahedron*. 1999;55(16):4999–5016.
 45. Yan P, Liu Y. Proline-Based 2,5-Diketopiperazine Alkaloids from Starfish *Pentacaster chinensis*. In: 2011 5th International Conference on Bioinformatics and Biomedical Engineering. 2011. p. 1–4.
 46. Palagiano E, De Marino S, Minale L, Riccio R, Zollo F, Iorizzi M, et al. Ptilomycalin A, crambescidin 800 and related new highly cytotoxic guanidine alkaloids from the starfishes *Fromia monilis* and *Celerina heffernani*. *Tetrahedron*. 1995;51(12):3675–82.
 47. Khadem S, Marles RJ. Chromone and flavonoid alkaloids: Occurrence and bioactivity. Vol. 17, *Molecules*. 2012. p. 191–206.
 48. Prompanya C, Fernandes C, Cravo S, Pinto MMM, Dethoup T, Silva AMS, et al. A new cyclic hexapeptide and a new isocoumarin derivative from the marine sponge-associated fungus *Aspergillus similanensis* KUFA 0013. *Mar Drugs*. 2015;13(3):1432–50.
 49. Sun K, Li Y, Guo L, Wang Y, Liu P, Zhu W. Indole diterpenoids and isocoumarin from the fungus, *Aspergillus flavus*, isolated from the prawn, *Penaeus vannamei*. *Mar Drugs*. 2014;12(7):3970–81.
 50. Kim DC, Quang TH, Ngan NTT, Yoon CS, Sohn JH, Yim JH, et al. Dihydroisocoumarin Derivatives from Marine-Derived Fungal Isolates and Their Anti-inflammatory Effects in Lipopolysaccharide-Induced BV2 Microglia. *J Nat Prod*. 2015;78(12):2948–55.
 51. Pan K, Tanaka C, Inagaki M, Higuchi R, Miyamoto T. Isolation and structure elucidation of GM4-type gangliosides from the Okinawan starfish *Protoreaster nodosus*. *Mar Drugs*. 2012;10(11):2467–80.
 52. Arunpanichler J, Rukachaisirikul V, Phongpaichit S, Supaphon O, Sakayaroj J. Meroterpenoid, isocoumarin, and phenol derivatives from the seagrass-derived fungus *Pestalotiopsis* sp. PSU-ES194. *Tetrahedron*. 2015;71(5):882–8.
 53. Berenbaum MR. The chemistry of defense: theory and practice. *Proc Natl Acad Sci* [Internet]. 1995;92(1):2–8. Available from: <http://www.pnas.org/cgi/doi/10.1073/pnas.92.1.2>
 54. Zagalsky PF, Haxo F, Hertzberg S, Hertzberg S, Llaaen-Jensen S. Studies on a blue carotenoprotein, linckiacyanin, isolated from the starfish *Linckia laevigata* (Echinodermata: Asteroidea). *Comp Biochem Physiol -- Part B Biochem*. 1989;93(2):339–53.
 55. Bandaranayake WM. The nature and role of pigments of marine invertebrates. Vol. 23, *Natural Product Reports*. 2006. p. 223–55.
 56. Mamelona J, Pelletier É, Girard-Lalancette K, Legault J, Karboune S, Kermasha S. Quantification of phenolic contents and antioxidant capacity of Atlantic sea cucumber, *Cucumaria frondosa*. *Food Chem* [Internet]. 2007 Jan;104(3):1040–7. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0308814607000684>
 57. Althunibat OY, Hashim R Bin, Taher M, Mohd. Daud J, Ikeda M-A, Zali BI. In Vitro Antioxidant and Antiproliferative Activities of Three Malaysian Sea Cucumber Species. *Eur J Sci Res*. 2009;37(3):376–87.
 58. Mamelona J, Pelletier É, Girard-Lalancette K, Legault J, Karboune S, Kermasha S. Antioxidants in digestive tracts and gonads of green urchin (*Strongylocentrotus droebachiensis*). *J Food Compos Anal*. 2011;24(2):179–83.
 59. Marmouzi I, Tamsouri N, El Hamdani M, Attar A, Kharbach M, Alami R, et al. Pharmacological and chemical properties of some marine echinoderms. *Rev Bras Farmacogn* [Internet]. 2018; Available from: <http://www.sciencedirect.com/science/article/pii/S0102695X17306853>
 60. Baharara J, Amini E. Phytochemical Screening, Antioxidant Effect and Down Regulation of TGF- β Induced by *Ophiocoma erinaceus* Brittle Star Crude Extract. *Zahedan J Res Med Sci* [Internet]. 2015 Dec 26;17(12). Available from: <http://zjrms.com/en/articles/5194.html>
 61. Lee CC, Hsieh HJ, Hsieh CH, Hwang DF. Antioxidative and anticancer activities of various ethanolic extract fractions from crown-of-thorns starfish (*Acanthaster planci*). *Environ Toxicol Pharmacol* [Internet]. 2014;38(3):761–73. Available from: <http://dx.doi.org/10.1016/j.etap.2014.08.021>
 62. Francis G, Kerem Z, Makkar HPS, Becker K. The biological action of saponins in animal systems: a review. *Br J Nutr* [Internet]. 2002;88(06):587. Available from: http://www.journals.cambridge.org/abstract_S0007114502002349
 63. Vincken JP, Heng L, de Groot A, Gruppen H. Saponins, classification and occurrence in the plant kingdom. *Phytochemistry*. 2007;68(3):275–97.
 64. Demeyer M, De Winter J, Caulier G, Eeckhaut I, Flammang P, Gerbaux P. Molecular diversity and body distribution of saponins in the sea star *Asterias rubens* by mass spectrometry. *Comp Biochem Physiol - B Biochem Mol Biol* [Internet]. 2014;168(1):1–11. Available from: <http://dx.doi.org/10.1016/j.cbpb.2013.10.004>
 65. Bahrami Y, Franco MC. Structure Elucidation of New Acetylated Saponins, Lessoniosides A, B, C, D, and E, and Non-Acetylated Saponins, Lessoniosides F and G, from the Viscera of the Sea Cucumber *Holothuria lessona*. Vol. 13, *Marine Drugs*. 2015.
 66. Yu S, Ye X, Chen L, Xie X, Zhou Q, Lian XY, et al. Cytotoxic and anti-colorectal tumor effects of sulfated saponins from sea cucumber *Holothuria moebii*. *Phytomedicine*. 2015;22(12):1112–9.
 67. Minale L, Riccio R, Zollo F. Steroidal oligoglycosides and polyhydroxysteroids from echinoderms. In: *Fortschritte der Chemie organischer Naturstoffe/Progress in the Chemistry of Organic Natural Products*. Springer; 1993. p. 75–308.
 68. De Marino S, Borbone N, McClintock JB, Amato F, Amato J, Amato J, et al. Bioactive Asterosaponins from the Starfish *Luidiaquinaria* and *Psilastercassiope*. Isolation and Structure Characterization by Two-Dimensional NMR Spectroscopy. *J Nat Prod* [Internet]. 2003;66(4):515–9. Available from: <http://pubs.acs.org/doi/abs/10.1021/np2005046>
 69. Tang HF, Yi YH, Li L, Sun P, Zhang SQ, Zhao YP. Asterosaponins from the starfish *Culcita novaeguineae* and their bioactivities. *Fitoterapia*. 2006;77(1):28–34.
 70. Malyarenko T V., Kicha AA, Ivanchina N V., Kalinovsky AI, Popov RS, Vishchuk OS, et al. Asterosaponins from the Far Eastern starfish *Leptasterias ochotensis* and their anticancer activity. *Steroids*. 2014;87:119–27.
 71. Thao NP, Cuong NX, Luyen BTT, Thanh N Van, Nhiem NX, Koh Y-S, et al. Anti-inflammatory Asterosaponins from the Starfish *Astropecten monacanthus*. *J Nat Prod* [Internet]. 2013 Sep 27;76(9):1764–70. Available from: <https://pubs.acs.org/doi/10.1021/np400492a>
 72. Riccio R, Greco OS, Minale L, Pusset J, Menou JL. Starfish saponins, part 18.1 steroidal glycoside sulfates from the starfish *Linckia laevigata*. *J Nat Prod*. 1985;48(1):97–101.

73. Minale L, Pizza C, Riccio R, Zollo F, Pusset J, Laboute P. Starfish Saponins, XIII. Occurrence of Nodososide in the Starfish *Acanthaster planci* and *Linckia laevigata* 1. J Nat Prod. 1984;47(3):558.
74. Riccio R, Zollo F, Finamore E, Minale L, Laurent D, Bargibant G, et al. Starfish saponins, 19. A novel steroidal glycoside sulfate from the starfishes protoreaster nodosus and pentaceraster alveolatus. J Nat Prod. 1985;48(2):266–72.
75. Thao NP, Thi T, Hanh H, Tu VA, Thi P, Huong M, et al. Chemical constituents from the starfish Protoreaster nodosus. Vietnam J Chem. 2015;53(2e):33–6.
76. Prabhu K, Bragadeeswaran S. Biological properties of brittle star *Ophiocnemis marmorata* collected from Parangipettai, Southeast coast of India. J Microbiol Antimicrob Full. 2013;5(October):110–8.
77. Mahato SB, Nandy AK, Roy G. Triterpenoids. Phytochemistry. 1992;31(7):2199–249.
78. Kalinin VI, Anisimov MM, Prokofieva NG, Avilov SA, Afiyatullof SS, Stonik VA. Biological activities and biological role of triterpene glycosides from holothurids (Echinodermata). Echinoderm Stud. 1996;5:139–81.
79. Kobayashi M, Okamoto Y, Kitagawa I. Marine natural products. XXVIII. The structures of sarasinosides A1, A2, A3, B1, B2, B3, C1, C2, and C3, nine new norlanostane-triterpenoidal oligoglycosides from the Palauan marine sponge *Asteropus sarasinosum*. Chem Pharm Bull. 1991;39(11):2867–77.
80. Li YX, Himaya SWA, Kim SK. Triterpenoids of marine origin as anti-cancer agents. Vol. 18, Molecules. 2013. p. 7886–909.
81. Jin DJ, Tang SA, Xing GS, Zhao WJ, Zhao C, Duan HQ, et al. Jaspiferins C-F, four new isomalabaricane-type triterpenoids from the South China Sea sponge *Jaspis stellifera*. J Asian Nat Prod Res. 2014;16(5):427–33.
82. Maier MS. Biological activities of sulfated glycosides from echinoderms. Stud Nat Prod Chem. 2008;35(C):311–54.
83. Patel SS, Savjani JK. Systematic review of plant steroids as potential anti-inflammatory agents: Current status and future perspectives. J Phytopharm [Internet]. 2015;4(42):121–5. Available from: www.phytopharmajournal.com
84. Kicha AA, Ivanchina N V., Kalinovskiy AI, Dmitrenok PS, Stonik VA. Sulfated steroid compounds from the starfish *Aphelasterias japonica* of the Kuril population. Russ Chem Bull. 2001;50(4):724–7.
85. Prokofieva NG, Chaikina EL, Kicha AA, Ivanchina N V. Biological activities of steroid glycosides from starfish. Comp Biochem Physiol - B Biochem Mol Biol. 2003;134(4):695–701.
86. Zhang W, Guo Y, Gu Y. Secondary Metabolites from the South China Sea Invertebrates: Chemistry and Biological Activity. 2006;2041–90.
87. D'Auria MV, Minale L, Riccio R. Polyoxygenated Steroids of Marine Origin. Chem Rev. 1993;93(5):1839–95.
88. Minale L, Pizza C, Riccio R, Zollo F. Steroidal glycosides from starfishes. Pure Appl Chem. 1982;54(10):1935–50.
89. Fleming WJ, Salathe R, Wyllie SG, Howden MEH. Isolation and partial characterization of steroid glycosides from the starfish *Acanthaster planci*. Comp Biochem Physiol Part B Comp Biochem [Internet]. 1976;53(3):267–72. Available from: <http://www.sciencedirect.com/science/article/pii/0305049176903230>
90. Itakura Y, Komori T. Biologically Active Glycosides from Asteroidea, X. Steroid Oligoglycosides from the Starfish *Acanthaster planci* L., 3. Structures of Four New Oligoglycoside Sulfates. Liebigs Ann der Chemie. 1986;1986(3):499–508.
91. Komori T, Matsuo J, Itakura Y, Sakamoto K, Ito Y, Taguchi S, et al. Biologically active glycosides from asteroidea, II. Steroid oligoglycosides from the starfish *Acanthaster planci* L., 1. Isolation and structure of the oligoglycoside sulfates. Liebigs Ann der Chemie. 1983;1983(1):24–36.
92. Qi J, Ojika M, Sakagami Y. linckosides A and B, two new neurotogenic steroid glycosides from the okinawan starfish *Linckia laevigata*. Bioorg Med Chem [Internet]. 2002;10(6):1961–6. Available from: <http://www.sciencedirect.com/science/article/pii/S0968089602000068>
93. Qi J, Ojika M, Sakagami Y. Linckosides C–E, three new neurotogenic steroid glycosides from the Okinawan starfish *Linckia laevigata*. Bioorg Med Chem [Internet]. 2004;12(15):4259–65. Available from: <http://www.sciencedirect.com/science/article/pii/S0968089604003670>
94. Calderón-Montaño J, Estefanía B-M, Manuel Luis O, Maldonado-Navas. Evaluating the Cancer Therapeutic Potential of Cardiac Glycosides. José Manuel. Biomed Res Int [Internet]. 2014;2014:1–9. Available from: <http://www.hindawi.com/journals/bmri/2014/794930/>
95. Felth J, Rickardson L, Rosén J, Wickström M, Fryknäs M, Lindskog M, et al. Cytotoxic effects of cardiac glycosides in colon cancer cells, alone and in combination with standard chemotherapeutic drugs. J Nat Prod. 2009;72(11):1969–74.
96. Newman R a, Yang P, Pawlus a D, Block KI. Cardiac glycosides as novel cancer therapeutic agents. MolInterv [Internet]. 2008;8:36–49. Available from: [y:5C%5CPUBLICATIONS%5CNEWMAN%5Cn2008%5Cnglycosides.pdf](http://www.scribd.com/document/100000000/y:5C%5CPUBLICATIONS%5CNEWMAN%5Cn2008%5Cnglycosides.pdf)
97. Kizer KW. Marine envenomations. Clin Toxicol. 1983;21(4–5):527–55.
98. Tlougan BE, Podjasek JO, Adams BB. Aquatic sports dermatoses. Part 2 - In the water: Saltwater dermatoses. Vol. 49, International Journal of Dermatology. 2010. p. 994–1002.

Cite this article: Walag AMP, Del Rosario R, Canencia OP. Zoochemical Composition of Selected Sea Stars Collected from the Coastal Waters of Carmen, Agusan Del Norte, Philippines. Asian J Biol Life Sci. 2019;8(2):53-62.